

In this response, we are not commenting on caveats such as aerosols, other greenhouse gases, lags, feedbacks, ice sheets, etc. The topic of *Shaviv and Veizer* [2003] was the “primary” climate driver on Phanerozoic time scales, with no space, or need, for any more discussion than that. Furthermore, we fail to see how any of the above would make CO₂ the “driver” in the Antarctic cores, when the temperature rises preceded those of CO₂ by centuries. We not only never denied but specifically highlighted the qualifying proposition that CO₂ may act as an amplifier.

In conclusion, the above response demonstrates that the “critique” of *Rahmstorf et al.* [2004] has little substance, in addition to the fact that it deals with time scales that are not even discussed in *Shaviv and Veizer* [2003]. Moreover, the statistical argument advanced in this issue of *Eos* as disproving the validity of the CRF/paleotemperature correlation is simply invalid (for details, see <http://www.phys.huji.ac.il/~shaviv/ClimateDebate/>).

References

Bond, G., B. Kromer, J. Beer, R. Muscheler, M. N. Evans, W. Showers, S. Hoffmann, R. Lotti-Bond, I. Hajdas, and G. Bonani (2001), Persistent solar influence

on North Atlantic climate during the Holocene, *Science*, 294(5549), 2130–2136.
 Boucot, A. J., and J. Gray (2001), A critique of Phanerozoic climatic modes involving changes in the CO₂ content of the atmosphere, *Earth Sci. Rev.*, 56(1-4), 1–159.
 Carslaw, K. S., R. G. Harrison, and J. Kirkby (2002), Cosmic rays, clouds and climate, *Science*, 298(5599), 1732–1737.
 Foukal, P. (2002), A comparison of variable solar total and ultraviolet irradiance outputs in the 20th century, *Geophys. Res. Lett.*, 29(23), 411–414, 2089, doi:10.2912002GL015474.
 Laut, P. (2003), Solar activity and terrestrial climate: An analysis of some purported correlations, *J. Atmos. Solar-Terr. Phys.*, 65(7), 801–812.
 Marsh, N. D., and H. Svensmark (2003), Galactic cosmic ray and El Niño–Southern Oscillation trends in ISCCP-D2 low cloud properties, *J. Geophys. Res.*, 108(D6), 4195, doi:10.1029/2001JD001264.
 Neff, U., S. J. Burns, A. Mangini, M. Mudelsee, D. Fleitmann, and A. Matter (2001), Strong coherence between solar variability and the monsoon in Oman between 9 and 6 kyr ago, *Nature*, 411(6835), 290–293.
 Rahmstorf, S., et al. (2004), Cosmic rays, carbon dioxide and climate, *Eos, Trans. AGU*, 85(4), 38–41.
 Rind, D. (2002), The Sun's role in climate variations, *Science*, 296(5568), 673–677.
 Royer, D. L., R. A. Berner, I. P. Montañez, N. J. Tabor, and D. J. Beerling (2004), CO₂ as a primary driver of Phanerozoic climate, *GSA Today*, 14(3), 4–10.

Shaviv, N. J. (2002), Cosmic ray diffusion from the galactic spiral arms, iron meteorites, and a possible climatic connection?, *Phys. Rev. Lett.*, 89(5), 051102.
 Shaviv, N. J. (2003), The spiral structure of the Milky Way, cosmic rays, and ice age epochs on Earth, *New Astron.*, 8(1), 39–77.
 Shaviv, N. J., and J. Veizer (2003), Celestial driver of Phanerozoic climate?, *GSA Today*, 13(7), 4–10.
 Solanki, S. K. (2002), Solar variability and climate change: is there a link?, *Astron. Geophys.*, 43(5), 5.9–5.13.
 Usoskin, I. G., S. K. Solanki, M. Schüssler, K. Mursula, and K. Alanko (2003), A millenium scale sunspot number reconstruction: evidence for an unusually Active Sun since the 1940's, *Phys. Rev. Lett.*, 91(21), 211101-1-211101-4.
 Veizer, J., et al. (1999), ⁸⁷Sr/⁸⁶Sr, δ¹³C and δ¹⁸O evolution of Phanerozoic seawater, *Chem. Geol.*, 161(1-3), 59–88.
 Veizer, J., Y. Godderis, and L. M. François (2000), Evidence for decoupling of atmospheric CO₂ and global climate during the Phanerozoic eon, *Nature*, 408(6813), 698–701.

—NIR SHAVIV, Racah Institute of Physics, Hebrew University of Jerusalem, Israel; and JAN VEIZER, Institut für Geologie, Mineralogie und Geophysik, Bochum, Germany, and Ottawa-Carleton Geoscience Centre, University of Ottawa, Ontario, Canada

Reply

PAGE 511

In our analysis [*Rahmstorf et al.*, 2004], we arrived at two main conclusions: the data of *Shaviv and Veizer* [2003] do not show a significant correlation of cosmic ray flux (CRF) and climate, and the authors' estimate of climate sensitivity to CO₂ based on a simple regression analysis is questionable. After careful consideration of Shaviv and Veizer's comment, we want to uphold and reaffirm these conclusions.

Concerning the question of correlation, we pointed out that a correlation arose only after several adjustments to the data, including shifting one of the four CRF peaks and stretching the time scale. To calculate statistical significance, we first need to compute the number of independent data points in the CRF and temperature curves being correlated, accounting for their autocorrelation. A standard estimate [*Quenouille*, 1952] of the number of effective data points is

$$N_{EFF} \equiv \frac{N}{1 + 2 \sum_{k=1}^N r_1(k) r_2(k)}$$

where N is the total number of data points and r_1, r_2 are the autocorrelations of the two series. For the curves of *Shaviv and Veizer* [2003], the result is $N_{EFF} = 4.8$. This is consistent with the fact that these are smooth curves with four humps, and with the fact that for CRF, the position of the four peaks is determined by four spiral arm crossings or four meteorite clusters, respectively; that is, by four independent data points. The number of points that enter the calculation of statistical significance of a

linear correlation is $(N_{EFF} - 2)$, since any curves based on only two points show perfect correlation; at least three independent points are needed for a meaningful result.

Shifting one of the four peaks to fit climate data reduces the number of independent points by one, and tuning the time scale to improve the fit uses up another degree of freedom, leaving between zero and one independent points in the significance calculation. Hence, no correlation is significant after the tuning steps of *Shaviv and Veizer* [2003]; given the few degrees of freedom in the data, the data were over-tuned. The fact that their tuning is within data uncertainty is irrelevant to statistical significance. It just means that a correlation might be possible without contradicting the data.

The consistency of the periods presented is still not convincing, since these periods are only averages of a few points with high variability. While it is possible that better data will demonstrate a correlation of cosmic rays and climate, our conclusion is that the data presented by *Shaviv and Veizer* [2003] are insufficient for this. As an aside, we did not confuse the exposure ages and real ages of meteorites.

Concerning the regression analysis to estimate climate sensitivity, Shaviv and Veizer write in their Comment, “we are not going to comment on caveats such as aerosols, other greenhouse gases, lags, feedbacks, ice sheets, etc.” This is unfortunate, since these issues are not caveats, but central to the determination of climate sensitivity to CO₂. As we pointed out, the strength of any individual forcing factor can only be estimated by a regression analysis if it is statistically independent from other forcings, which is very unlikely for the examples mentioned, or if these other forcings are explicitly taken

into account, as in *Lorius et al.* [1990]. Since this was not done, we maintain that the regression is questionable.

Finally, it is worth pointing out areas of agreement.

Shaviv and Veizer state, “we fail to see how any of the above would make CO₂ the ‘driver’ in the Antarctic ice cores.” We fully agree that CO₂ is not the driver of the climate variability seen in these cores. There is a host of excellent empirical evidence and widespread agreement that climate variability on glacial-interglacial time scales is driven by variations in the Earth's orbit, the Milankovich cycles, with CO₂ responding as a positive feedback.

The earliest analysis of Antarctic cores, and the derivation of climate sensitivity from these data, was already based on this premise (see *Lorius et al.* [1990]). Hence, climatologists have long expected a time lag of CO₂ behind temperature in the ice core data, and some of us were involved in pioneering the measurement of this lag using a gas-based temperature proxy that resolves the problem of the age difference between gas bubbles and the surrounding ice [*Caillon et al.*, 2003]. The result is a lag of 800 years at termination III (240,000 yr B.P.), a warming that occurred over a 5000-yr period.

This means that one-sixth of the warming at the end of this glacial period occurred before the CO₂ feedback started to be felt. This is consistent with recent climate model simulations of glacial cycles, which show that CO₂ changes are not required to explain the initiation of glaciation or deglaciation, but that the CO₂ feedback is needed to explain their full extent [*Yoshimori et al.*, 2001; *Meissner et al.*, 2003]. The time lag in ice core data gives no information about the climate sensitivity to a given CO₂ change, such as that caused by anthropogenic emissions.

We also fully agree with Shaviv and Veizer that their results, even if they were correct, apply only to the multi-million-year time scale and cannot be applied to shorter time scales. We are glad they have clarified this point. Their media releases as well as their paper, in which they compare their climate sensitivity with the range given by the Intergovernmental Panel on Climate Change (which applies to modern climate and centennial time scales [IPCC, 2001]), could have been misunderstood in this respect.

References

Caillon, N., J. P. Severinghaus, J. Jouzel, J.-M. Barnola, J. Kang, and V. Y. Lipenkov (2003), Timing of atmospheric CO₂ and Antarctic temperature changes across termination III, *Science*, 299, 1728–1731.

Intergovernmental Panel on Climate Change (2001), *Climate Change 2001*, Cambridge University Press, Cambridge.
 Lorius, C., J. Jouzel, D. Raynaud, J. Hansen, and H. Le Treut (1990), The ice-core record: Climate sensitivity and future greenhouse warming, *Nature*, 347 (13 Sept. 1990), 139–145.
 Meissner, K. J., A. J. Weaver, H. D. Matthews, and P. M. Cox (2003), The role of land-surface dynamics in glacial inception: A study with the UVic Earth System Model, *Clim. Dyn.*, 21 (7–8), 515–537.
 Quenouille, M. H. (1952), *Associated Measurements*, 242 pp., Butterworth Scientific Publications, London.
 Rahmstorf, S., et al. (2004), Cosmic rays, carbon dioxide and climate, *Eos, Trans. AGU*, 85(4), 38, 41.
 Shaviv, N., and J. Veizer (2003), Celestial driver of Phanerozoic climate?, *GSA Today*, 13(7), 4–10.
 Yoshimori, M., A. J. Weaver, S. J. Marshall, and G. K. C. Clarke (2001), Glacial termination: Sensitivity to orbital and CO₂ forcing in a coupled climate system model, *Clim. Dyn.*, 17 (8), 571–588.

—STEFAN RAHMSTORF, Potsdam Institute for Climate Impact Research, Germany; DAVID ARCHER, University of Chicago, Ill.; DENTON S. EBEL, American Museum of Natural History, N.Y.; OTTO EUGSTER, Dept. of Space Research and Planetology, Bern, Switzerland; JEAN JOUZEL, Institut Pierre Simon Laplace/LSC, Saclay, France; DOUGLAS MARAUN, Institute of Physics, Potsdam University, Germany; URS NEU, ProClim, Swiss Academy of Sciences, Bern, Switzerland; GAVIN A. SCHMIDT, NASA Goddard Institute for Space Studies and Center for Climate Systems Research, Columbia University, N.Y.; JEFF SEVERINGHAUS, Scripps Institution of Oceanography, San Diego, Calif.; ANDREW J. WEAVER, School of Earth and Ocean Sciences, University of Victoria, Canada; and JIM ZACHOS, University of California, Santa Cruz

SECTION NEWS

SPACE PHYSICS & AERONOMY




Section President, Michael J. Mendillo;
Section Secretaries, Gang Lu, William S. Kurth, and Michelle F. Thomsen

Murr Receives 2004 F. L. Scarf Award

PAGE 511

David Murr has been awarded the F.L. Scarf Award given annually to a recent Ph.D. recipient for outstanding dissertation research that contributes directly to solar-planetary sciences. Murr's thesis is entitled "Magnetosphere-iono-



sphere coupling on meso- and macro-scales." He will be formally presented with the award on 14 December during the 2004 AGU Fall Meeting,

which is held 13–17 December in San Francisco, California.

Murr received his B.S. in physics from Augsburg College, Minneapolis, Minnesota, in 1992 under the direction of Mark Engebretson. After spending two years in the Peace Corps and an additional two years working in industry, he returned to the field of space physics as project manager for the MACCS (Magnetometer Array for Cusp and Cleft Studies) array of magnetometers at Boston University. Supervised by W. J. Hughes, he earned his Ph.D. at Boston University in 2003. Also in 2003, he received the National Science Foundation Geospace Environment Modeling (GEM) postdoctoral researcher award.

Murr currently works with William Lotko at Dartmouth College, Hanover, New Hampshire, studying the transient response of the dayside magnetosphere to rapid solar wind forcing and the outflow of ionospheric plasma into the magnetosphere.

BOOK REVIEW

Measuring the Oceans from Space



IAN S. ROBINSON

Springer-Praxis; New York; ISBN 3-540-42647-7; xlv+669 pp., 23 color plates; 2004; \$189.

PAGE 514

Given the growth in the number and complexity of ocean satellite systems since the launch of Seasat in 1978, combined with the fact that about twenty-five countries are now involved in such observations, this is an appropriate time for a new book on satellite

oceanography. (Disclosure: the book reviewer has also just published an ocean remote sensing book.) The present book is an expanded version of Robinson's 1985 book, *Satellite Oceanography*.

It consists of two parts: fundamentals of satellite oceanography and remote sensing techniques. Although Robinson originally planned to include a third part on ocean applications, because of the length of the current book, that section will be published separately in 2005. The present book discusses for the ice-free ocean "what is measured and how it is done."

The five chapters in Part I occupy about a quarter of the text. The topics covered include a history of satellite oceanography, the

electromagnetic spectrum, the different kinds of sensors, and a qualitative introduction to atmospheric radiative transfer. Topics also include platforms, orbits and their scales of temporal and spatial coverage, data encoding, and image processing.

Part II contains seven chapters: six on the various instruments, and a conclusion. The instrument chapters cover the visible (ocean color), infrared sea surface temperature (SST), passive microwave, scatterometry, synthetic aperture radar (SAR), and altimetry.

Each chapter discusses the form of the atmospheric radiative transfer equation appropriate to the instrument, the interaction of this radiation with the ocean surface and interior, and how the instrument retrieves the property in question. The chapters next consider past, present, and near-future instruments, with particular emphasis on European instruments, continue with applications, and