



The 8.2 ka event: Abrupt transition of the subpolar gyre toward a modern North Atlantic circulation

A. Born

Bjerknes Centre for Climate Research, Allégaten 55, N-5007 Bergen, Norway (andreas.born@bjerknes.uib.no)

Also at Geophysical Institute, University of Bergen, N-5007 Bergen, Norway

A. Levermann

Potsdam Institute for Climate Impact Research, D-14412 Potsdam, Germany

Also at Institute of Physics, Potsdam University, D-14469 Potsdam, Germany

[1] Climate model simulations of the 8.2 ka event show an abrupt strengthening of the Atlantic subpolar gyre that allows us to connect two major but apparently contradictory climate events of the early Holocene: the freshwater outburst from proglacial lakes and the onset of Labrador Sea water formation. The 8.2 ka event is the largest climatic signal of our present interglacial with a widespread cooling in the North Atlantic region about 8200 years before present. It coincides with a meltwater outburst from North American proglacial lakes that is believed to have weakened the Atlantic meridional overturning circulation and northward heat transport, followed by a recovery of the deep ocean circulation and rising temperatures after a few centuries. Marine proxy data, however, date the onset of deep water formation in Labrador Sea to the same time. The subsequent strengthening of the slope current system created a regional signal recorded as an abrupt and persistent surface temperature decrease. Although similarities in timing are compelling, a mechanism to reconcile these apparently contradictory events was missing. Our simulations show that an abrupt and persistent strengthening of the Atlantic subpolar gyre provides a plausible explanation. The intense freshwater pulse triggered a transition of the gyre circulation into a different mode of operation, stabilized by internal feedbacks and persistent after the cessation of the perturbation. As a direct consequence, deep water formation around its center intensifies. This corresponds to the modern flow regime and stabilizes the meridional overturning circulation, possibly contributing to the Holocene's climatic stability.

Components: 5100 words, 5 figures.

Keywords: 8.2 ka event; Holocene; North Atlantic; climate model; subpolar gyre.

Index Terms: 4901 Paleoceanography: Abrupt/rapid climate change (1605); 0473 Biogeosciences: Paleoclimatology and paleoceanography (3344); 4936 Paleoceanography: Interglacial.

Received 4 January 2010; **Revised** 10 May 2010; **Accepted** 17 May 2010; **Published** 25 June 2010.

Born, A., and A. Levermann (2010), The 8.2 ka event: Abrupt transition of the subpolar gyre toward a modern North Atlantic circulation, *Geochem. Geophys. Geosyst.*, 11, Q06011, doi:10.1029/2009GC003024.

1. Introduction

[2] During the relatively stable conditions of our present interglacial, the 8.2 ka event is the largest climatic signal with a widespread cooling in the North Atlantic region about 8200 years before present. It coincides with a meltwater outburst from North American proglacial lakes [Alley *et al.*, 1997; Alley and Augústsdóttir, 2005, and references therein]. In current understanding, this caused a weakening of the Atlantic meridional overturning circulation (AMOC) and a subsequent reduction in northward heat transport, followed by a recovery of the deep ocean circulation and rising temperatures after a few centuries [Bauer *et al.*, 2004; Hall *et al.*, 2004; Ellison *et al.*, 2006; Wiersma *et al.*, 2006; Kleiven *et al.*, 2008].

[3] This two-dimensional explanation, however, is not able to explain a number of marine paleorecords that clearly call for a three-dimensional mechanism. An abrupt but persistent surface temperature decrease was reported from the western North Atlantic at the time of the 8.2 ka event (Figure 1) [Solignac *et al.*, 2004; de Vernal and Hillaire-Marcel, 2006; Sachs, 2007]. It has been suggested to be associated with the onset of deep water formation in Labrador Sea and a subsequent strengthening of the slope current system, the western branch of the subpolar gyre (SPG) [Hillaire-Marcel *et al.*, 2001; Sachs, 2007]. Proxy data from Reykjanes Ridge shows a similarly abrupt and persistent warming which provides evidence for an enhanced Irminger Current (Figure 1) [Andersen *et al.*, 2004; Came *et al.*, 2007]. This is the northeastern branch of the SPG and thus corroborates the hypothesis of a stronger gyre and enhanced convection in its center.

[4] The concurrence of these two events raises the question of how convection can increase at the time of the most severe freshwater flood of the past 10,000 years. We argue, based on coupled climate model experiments, that this is no contradiction but that a causal relationship connects the two events. A change in the density structure is found to provide positive feedbacks leading to a permanent strengthening of the SPG and convection. The impact of baroclinic adjustments on the SPG strength is well documented [Eden and Willebrand, 2001; Häkkinen and Rhines, 2004; Hátún *et al.*, 2005; Treguier *et al.*, 2005; Levermann and Born, 2007; Born *et al.*, 2009; Lohmann *et al.*, 2009; Born *et al.*, 2010; Montoya *et al.*, 2010] and this

study aims to apply this physical understanding to data from the geological record.

2. Model Description

[5] To investigate the role of the SPG during the 8.2 ka event we make use of the coupled climate model CLIMBER-3 α , which comprises atmosphere and sea ice components and the oceanic general circulation model MOM-3 (Montoya *et al.* [2005] and auxiliary material).¹ The oceanic horizontal resolution is $3.75^\circ \times 3.75^\circ$ with 24 unevenly spaced vertical layers. Simplified atmosphere dynamics are solved on a coarse numerical grid (7.5° in latitude and 22.5° in longitude).

[6] Due to the coarse resolution of the model, the SPG and deep water formation in its center are shifted eastward and simulated boundary currents are relatively broad (Figure 2). A second deep water formation region is found in the Nordic seas. Despite shortcomings, the interaction between gyre advection, convection in its center and eddy transport in between, as well as the communication of deep water masses across the deepest passages of the Greenland-Scotland ridge, compares well to higher-resolution models [Spall, 2005; A. Born *et al.*, Late Eemian warming in the Nordic seas as seen in proxy data and climate models, submitted to *Paleoceanography*, 2010]. For additional information we refer to sensitivity studies with glacial [Montoya *et al.*, 2010] and present-day boundary conditions [Mignot *et al.*, 2006] as well as under global warming [Levermann *et al.*, 2007].

[7] For the present study, the model was initialized with climatological hydrography [Levitus, 1982] and orbital parameters for 8,200 years before present [Berger, 1978]. Atmospheric CO₂ concentration was set to 260 ppm [Raynaud *et al.*, 2000]. The continental watershed over North America was shifted to the west and south in order to take into account changes in surface gradient due to the isostatic depression of glacial ice sheets. After the model was run into equilibrium over 2700 years, the simulated SPG exhibits a stable but weak volume transport of 19.5 Sv (1 Sv = 10^6 m³ s⁻¹). In order to simulate the lake drainage, $160 \cdot 10^{12}$ m³ of freshwater were added to the surface of the Labrador Sea coast during a period of two years, west of the deep convection region. This value is equivalent to a 2 year volume

¹Auxiliary materials are available in the HTML. doi:10.1029/2009GC003024.




Figure 1. Map of the subpolar North Atlantic showing the location of core sites and of major ocean currents mentioned in the text. Blue dots denote cores that show an abrupt and persistent cooling after 8 ka before present; warming is reported from the Reykjanes Ridge and marked red. Grey dots show where the 8.2 ka event is evident as a temporal reduction in deep current flow speed. Black arrows illustrate the surface currents (IC, Irminger Current; EGC, East Greenland Current; LC, Labrador Current; NAC, North Atlantic Current); grey arrows illustrate the Deep Western Boundary Current. For a list of full core names and references, please refer to Table S1 in Text S1.

flux of 2.6 Sv, based on a recent estimate of the lake volume [Leverington *et al.*, 2002], and has been used in previous model studies [Bauer *et al.*, 2004; LeGrande *et al.*, 2006; Wiersma *et al.*, 2006]. For diagnostic purposes, a passive tracer was released simultaneously with the freshwater flux at the same location in order to track advection of the perturbation. This tracer does not influence the circulation. We define the strength of the subpolar gyre as the local minimum of the depth integrated stream function. For the cyclonic circulation of the SPG, this stream function is negative (Figure 2).

3. Model Results and Interpretation of Proxy Data

[8] In response to the meltwater release, the SPG switches into a significantly stronger mode with 29 Sv volume transport (Figures 2 and 3a). This represents a strengthening of about 50% and the stronger circulation is within the uncertainty of present-day observations [Bacon, 1997; Read, 2001]. Further integration shows that this stronger state is stable and not only a temporary response to the freshwater pulse.

[9] The abrupt transition is due to two positive feedbacks inherent to the SPG [Levermann and Born, 2007]. First, a stronger SPG transports less tropical saline water into the Nordic seas but accu-

mulates these in the subpolar North Atlantic, making upper water masses in the center of the SPG more saline (Figures S7–S9 in Text S1). This is consistent with previous conclusions which are based on simulations with a high-resolution ocean general circulation model and confirmed by observations [Hátún *et al.*, 2005; Lohmann *et al.*, 2009]. Increased surface salinity enhances deep convection, cooling the deep water column. Besides, a stronger SPG results in enhanced outcropping of isopycnals




Figure 2. Colors indicate anomaly of the vertically integrated stream function, last 50 years of the perturbed simulation minus last 50 years before the meltwater pulse. Contours indicate vertically integrated stream function for the last 50 years of the simulation. For the cyclonic circulation of the SPG, the negative anomaly represents a strengthening. Outside the subpolar North Atlantic, currents are virtually unchanged.




Figure 3. Temporal evolution of key quantities of the transition toward a stronger SPG during the 8.2 ka event. The vertical line indicates the timing of the lake Agassiz drainage; data are filtered with a 25 year running mean. (a) Volume and heat transport of the SPG, (b) sea surface temperature in the northeastern (black) and southwestern (red) subpolar region, and (c) maximum winter mixed layer depth in the center of the SPG and surface density in the center of the SPG (see Figure S7 in Text S1). The stronger heat transport leads to a warming of the northeastern gyre region while the southwestern part cools rapidly. Surface density and mixed layer depth increase simultaneously.

and hence a more efficient removal of heat from the gyre's center by isopycnal mixing. Both effects, salt accumulation and removal of heat, increase the core density of the gyre compared to the relatively light exterior, sea surface drops and the corresponding geostrophic response strengthens the cyclonic SPG circulation.

[10] In addition to these self-sustaining internal feedbacks, there exists an interaction with the flow over the Greenland-Scotland ridge. The meltwater perturbation reduces sinking in the Nordic seas and therewith the supply of dense overflow waters to the northern rim of the SPG. Consequently, the outer rim of the gyre gets lighter and the gyre slightly intensifies. This triggers the two internal feedbacks mentioned above and yields the stronger SPG state

(Figure S1 in Text S1). Elsewhere, we have shown that a short reduction of dense deep outflow from the Nordic seas is sufficient to trigger the transition (≥ 25 years [Levermann and Born, 2007]) and that the described internal feedbacks dominate the dynamics. Changes in wind stress are small and do not show a consistent pattern (Figure S6 in Text S1). This is plausible because the freshwater flood does not impact wind patterns directly. Moreover, the coarse resolution of the atmosphere component might underestimate regional variations. Thus, the changes in wind stress seen after the transition are likely due to local changes in sea ice as suggested also by the irregular pattern. The mechanism of transition is robust to prescribed wind forcing and with respect to model setup and experimental design (see auxiliary material).

[11] Considering this change in surface circulation allows us to combine a number of proxy records into a consistent picture of the 8.2 ka event. Recent work found strong evidence for a reduction of dense water supply from the Nordic seas due to the lake Agassiz drainage [Hall et al., 2004; Ellison et al., 2006; Kleiven et al., 2008]. However, proxy data suggests that the meltwater signal did not reach the convection region at the center of the Labrador Sea but was exported along the shelf in the Labrador Current [Keigwin et al., 2005; Hillaire-Marcel et al., 2007]. From there a large portion moved northeastward and was diluted by mixing with water of the North Atlantic Current before reaching the Nordic seas and causing a moderate reduction in convection there. Our model reproduces this path qualitatively. 30 years after the lake drainage, highest concentrations of meltwater are found in the Nordic seas where 25% of the meltwater pulse has been advected to (Figure 4).

[12] After the transition, the stronger SPG circulation intensifies oceanic heat transport (Figure 3a). More warm tropical water reaches the northern SPG while more cold water is advected toward the west and south. This results in a sea surface temperature (SST) dipole (Figure S7 in Text S1). The southwestern SPG region cools abruptly (Figure 3b, red) while rapid warming is seen in the northeastern SPG region (Figure 3b, black).

[13] An abrupt and persistent SST decrease coeval with the lake drainage has indeed been reported from dinoflagellate cysts and alkenones for many locations throughout Labrador Sea and downstream Labrador Current south of Newfoundland [Solignac et al., 2004; de Vernal and Hillaire-Marcel, 2006; Sachs, 2007]. Consistent with our simulations this




Figure 4. Distribution of the meltwater pulse 30 years after its release into the Labrador Sea, integrated over depth (in m). Highest concentrations are found in the Nordic seas. Inset shows the fraction of the pulse in the Nordic seas and the Arctic Ocean. After approximately 30 years, 25% of the meltwater is advected north of the Greenland-Scotland ridge, followed by a decline as it is diluted globally.

data indicates a drastic reorganization of the slope current system. No persistent cooling is seen in foraminiferal data from Eirik Drift in the northern Labrador Sea, probably because of its location too far offshore to record changes in the coastal current [Kleiven *et al.*, 2008]. The warming signal on the eastern side of the gyre is not easily detected in proxy records because the strengthening of the surface circulation also changed the position of frontal systems. Thus, the strengthening of the SPG is recorded as a subsurface cooling inferred from foraminifera at a relatively eastern location [Thornalley *et al.*, 2009]. However, approximately 400 km further west on Reykjanes Ridge, where currents are more constrained by topography, an increase and stabilization of surface temperatures reconstructed using diatoms, foraminiferal $\delta^{18}\text{O}$ and Mg/Ca data has been associated with an intensification of the Irminger Current, the northern limb of the SPG (Figure 1) [Andersen *et al.*, 2004; Came *et al.*, 2007].

[14] Further support for a transition toward a stronger SPG following the 8.2 ka event comes from increased sea surface salinities throughout the subpolar region that started similarly abrupt as the temperature changes and were equally persistent, both in our model experiments (Figures S7–S9 in Text S1) and proxy data [Solignac *et al.*, 2004; de Vernal and Hillaire-Marcel, 2006; Hillaire-Marcel *et al.*, 2007; Came *et al.*, 2007]. The cooling of western slope waters can thus not be attributed to more intense transport of cold and fresh Arctic waters through the East Greenland Current. Their upstream source must be the rela-

tively saline Irminger Current which suggests a SPG intensification. A sudden increase in advection of Atlantic water into Labrador Sea has also been concluded from diatoms, fossil shell data and foraminiferal assemblages at several locations along the Greenland west coast [Donner and Jungner, 1975; Lloyd *et al.*, 2005; Ren *et al.*, 2009]. The intensification of westward salt transport and the eastward movement of the subpolar front with a stronger SPG results in a freshening of inflow into the Nordic seas (Figure S7 in Text S1), consistent with paleo-observations [Thornalley *et al.*, 2009] and present-day instrumental records [Hátún *et al.*, 2005]. Salt accumulation in the center of an anomalously strong SPG has been shown in a higher-resolution model [Lohmann *et al.*, 2009].

[15] Proxy data of Labrador Sea convection furthermore supports a significant reorganization of the surface circulation. In our simulations, enhanced surface salinity together with subsurface cooling in the center of the SPG results in a density increase and a subsequent intensification of convection south of the Greenland-Scotland ridge (Figure 3c). These results match well with existing data of the geological record. Little or no Labrador Sea Water was produced in the early Holocene and formation intensified not long after the meltwater outburst [Hillaire-Marcel *et al.*, 2001, 2007]. Isotopic changes in Labrador Sea sediments show a persistent reorganization of the deep current system after 8,000 years before present [Fagel *et al.*, 2004] with deeper Labrador Sea Water found in Pa/Th data [Gherardi *et al.*, 2009] and a permanent reduction of deep water formed in the Nordic seas




Figure 5. The Atlantic Meridional Overturning Circulation (AMOC), decomposed into deep water contributions from sinking in the Nordic seas (blue) and south of the Greenland-Scotland ridge (red). The vertical line indicates the timing of the lake Agassiz drainage, and a 25 year running mean filter is applied. The AMOC weakens abruptly in response to the freshwater event and recovers gradually over approximately 100 years. While the initial weakening is caused by the reduction of deep water supply from the Nordic seas, the recovery is due to a delayed intensification of Labrador Sea deep water formation. Both Nordic seas and Labrador Sea downwelling change significantly and persistently.

recorded in benthic $\delta^{13}\text{C}$ data [Evans *et al.*, 2007]. This points toward a persistent reorganization of both the horizontal surface and the deep meridional circulation, as simulated here.

[16] In contrast to significant changes in SPG strength (47%), the AMOC weakens by just 1.5 Sv (11%) in response to the meltwater pulse, significantly less than the 30%–50% reduction reported from previous model studies [Bauer *et al.*, 2004; Wiersma *et al.*, 2006; LeGrande *et al.*, 2006] (Figure 5). The strong response in earlier simulations might be due to the application of the meltwater pulse directly on the Labrador Sea convection region with dramatic consequences for SPG and AMOC. Such a freshwater perturbation leads to a reduction of convection in the Labrador Sea contradicting the paleorecord that shows the intensification of convection in this region. Moreover, this scenario removes a significant fraction of the freshwater pulse from the surface, inconsistent with proxy data indicating its advection in the Labrador Current. Hence, a smaller fraction is advected into the Nordic seas which is crucial to initiate the mechanism amplifying the SPG as discussed above (see auxiliary material for further discussion). The initial AMOC weakening in our simulation is followed by a rapid recovery over approximately 100 years and then a more gradual

increase. The decomposition into deep water formation regions shows that the initial weakening is due to reduced sinking in the Nordic seas, reaching its minimum after 150 years. The recovery and weak overall reduction is due to an increase in deep water formation south of the Greenland-Scotland ridge with a time lag of approximately 50 years (Figures 5 and 3c).

[17] A distinctive reduction in deep current flow speeds, probably delayed by at least several decades, is reported from the Greenland-Scotland ridge overflows and off the southern tip of Greenland (Figure 1) [Hall *et al.*, 2004; Ellison *et al.*, 2006; Kleiven *et al.*, 2008]. This is upstream from where Labrador Sea Water joins the Deep Western Boundary Current, the southward flowing branch of the AMOC. Further south in the North Atlantic, the reduction of northern source deep waters is minor for the 8.2 ka event if recorded at all [Keigwin and Boyle, 2000; Oppo *et al.*, 2003; Keigwin *et al.*, 2005]. The onset of Labrador Sea Water formation following the meltwater pulse might explain this discrepancy.

4. Summary and Conclusions

[18] Expanding previous studies that focused on the AMOC response, i.e., changes in the meridional circulation, to the lake Agassiz drainage, we propose that many observed but hitherto unexplained abrupt changes and discrepancies require taking into account a reorganization of the horizontal circulation. The most striking result is the persistent strengthening of the SPG in response to the short freshwater pulse. The transition between the two circulation patterns is triggered by an external positive feedback and stabilized by two positive feedback mechanisms within the SPG.

[19] While our results do not contradict an abrupt and considerable AMOC reduction in response to the lake Agassiz drainage, it might have been relatively short lived and weaker than suggested by previous simulations for two reasons: (1) The freshwater flood had the biggest impact not in the Labrador Sea but primarily affected deep water formation in the Nordic seas after mixing in the North Atlantic Current (Figure 4) and (2) this deep water reduction was partly compensated by enhanced sinking in the Labrador Sea (Figures 3c and 5). The latter mechanism has already been observed in models and data on time scales of several millennia [Renssen *et al.*, 2005; Solignac *et al.*, 2004].

[20] In our model the reorganization of the SPG surface circulation and subsequent changes in heat and salt advection provide the precondition for a more intense Labrador Sea convection and stabilize it. It has been suggested that this circulation mode is a unique feature of the Holocene, with Labrador Sea Water probably missing in the warmer climate of the last interglacial [Hillaire-Marcel *et al.*, 2001] and implications for the stability of the AMOC by the end of this century. Our results might provide the base for a future investigations of these hypotheses.

Acknowledgments

[21] We gratefully acknowledge useful discussions with Ulysses S. Ninnemann, Kerim H. Nisancioglu, and Trond Dokken. An earlier version of this manuscript greatly benefited from comments of three anonymous reviewers. A.B. was funded by the Marie Curie Actions project NICE (MRTN-CT-2006-036127). This is publication A290 from the Bjerknæs Centre for Climate Research.

References

- Alley, R. B., and A. M. Ágústssdóttir (2005), The 8k event: Cause and consequences of a major Holocene abrupt climate change, *Quat. Sci. Rev.*, *24*, 1123–1149.
- Alley, R. B., P. A. Mayewski, T. Sowers, M. Stuiver, K. C. Taylor, and P. U. Clark (1997), Holocene climatic instability: A prominent, widespread event 8200 years ago, *Geology*, *25*, 483–486.
- Andersen, C., N. Koc, and M. Moros (2004), A highly unstable Holocene climate in the subpolar North Atlantic: Evidence from diatoms, *Quat. Sci. Rev.*, *23*, 2155–2166.
- Bacon, S. (1997), Circulation and fluxes in the North Atlantic between Greenland and Ireland, *J. Phys. Oceanogr.*, *27*, 1420–1435.
- Bauer, E., A. Ganopolski, and M. Montoya (2004), Simulation of the cold climate event 8200 years ago by meltwater outburst from Lake Agassiz, *Paleoceanography*, *19*, PA3014, doi:10.1029/2004PA001030.
- Berger, A. (1978), Long-term variations of caloric solar radiation resulting from the Earth's orbital elements, *Quat. Res.*, *9*, 139–167.
- Born, A., A. Levermann, and J. Mignot (2009), Sensitivity of the Atlantic Ocean circulation to a hydraulic overflow parameterisation in a coarse resolution model: Response of the subpolar gyre, *Ocean Modell.*, *27*(3–4), 130–142.
- Born, A., K. H. Nisancioglu, and P. Braconnot (2010), Sea ice induced changes in ocean circulation during the Eemian, *Clim. Dyn.*, doi:10.1007/s00382-009-0709-2, in press.
- Came, R. E., D. W. Oppo, and J. F. McManus (2007), Amplitude and timing of temperature and salinity variability in the subpolar North Atlantic over the past 10 k.y., *Geology*, *35*, 315–318.
- de Vernal, A., and C. Hillaire-Marcel (2006), Provincialism in trends and high frequency changes in the northwest North Atlantic during the Holocene, *Global Planet. Change*, *54*, 263–290.
- Donner, J., and H. Jungner (1975), Radiocarbon dating of shells from marine deposits in the Disko Bugt, West Greenland, *Boreas*, *4*, 25–45.
- Eden, C., and J. Willebrand (2001), Mechanism of interannual to decadal variability of the North Atlantic Circulation, *J. Clim.*, *14*, 2266–2280.
- Ellison, C. R. W., M. R. Chapman, and I. R. Hall (2006), Surface and deep ocean interactions during the cold climate event 8200 years ago, *Science*, *312*, 1929–1932.
- Evans, H. K., I. R. Hall, G. G. Bianchi, and D. W. Oppo (2007), Intermediate water links to Deep Western Boundary Current variability in the subtropical NW Atlantic during marine isotope stages 5 and 4, *Paleoceanography*, *22*, PA3209, doi:10.1029/2006PA001409.
- Fagel, N., C. Hillaire-Marcel, M. Humblet, R. Brasseur, D. Weis, and R. Stevenson (2004), Nd and Pb isotope signatures of the clay-size fraction of Labrador Sea sediments during the Holocene: Implications for the inception of the modern deep circulation pattern, *Paleoceanography*, *19*, PA3002, doi:10.1029/2003PA000993.
- Gherardi, J.-M., L. Labeyrie, S. Nave, R. Francois, J. F. McManus, and E. Cortijo (2009), Glacial-interglacial circulation changes inferred from ²³¹Pa/²³⁰Th sedimentary record in the North Atlantic region, *Paleoceanography*, *24*, PA2204, doi:10.1029/2008PA001696.
- Häkkinen, S., and P. B. Rhines (2004), Decline of subpolar North Atlantic circulation during the 1990s, *Science*, *304*, 555–559.
- Hall, I. R., G. Bianchi, and J. R. Evans (2004), Centennial to millennial scale Holocene climate-deep water linkage in the North Atlantic, *Quat. Sci. Rev.*, *23*, 1529–1536.
- Hátún, H., A. B. Sandø, H. Drange, B. Hansen, and H. Valdimarsson (2005), Influence of the Atlantic Subpolar Gyre on the thermohaline circulation, *Science*, *309*, 1841–1844.
- Hillaire-Marcel, C., A. de Vernal, A. Bilodeau, and A. J. Weaver (2001), Absence of deep-water formation in the Labrador Sea during the last interglacial period, *Nature*, *410*, 1073–1077.
- Hillaire-Marcel, C., A. de Vernal, and D. J. W. Piper (2007), Lake Agassiz Final drainage event in the northwest North Atlantic, *Geophys. Res. Lett.*, *34*, L15601, doi:10.1029/2007GL030396.
- Keigwin, L. D., and E. A. Boyle (2000), Detecting Holocene changes in thermohaline circulation, *Proc. Natl. Acad. Sci. U. S. A.*, *97*, 1343–1346.
- Keigwin, L. D., J. P. Sachs, Y. Rosenthal, and E. A. Boyle (2005), The 8200 year B.P. event in the slope water system, western subpolar North Atlantic, *Paleoceanography*, *20*, PA2003, doi:10.1029/2004PA001074.
- Kleiven, H. F., C. Kissel, C. Laj, U. S. Ninnemann, T. O. Richter, and E. Cortijo (2008), Reduced North Atlantic Deep Water coeval with the glacial Lake Agassiz fresh water outburst, *Science*, *319*, 60–64.
- LeGrande, A. N., G. A. Schmidt, D. T. Shindell, C. V. Field, R. L. Miller, D. M. Koch, G. Faluvegi, and G. Hoffmann (2006), Consistent simulations of multiple proxy responses to an abrupt climate change event, *Proc. Natl. Acad. Sci. U. S. A.*, *103*, 837–842.
- Leverington, D., H. D. Mann, and J. T. Teller (2002), Changes in bathymetry and volume of glacial Lake Agassiz between 9200 and 7700 14C yr BP, *Quat. Res.*, *57*, 244–252.
- Levermann, A., and A. Born (2007), Bistability of the Atlantic subpolar gyre in a coarse-resolution climate model, *Geophys. Res. Lett.*, *34*, L24605, doi:10.1029/2007GL031732.

- Levermann, A., J. Mignot, S. Nawrath, and S. Rahmstorf (2007), The role of northern sea ice cover for the weakening of the thermohaline circulation under global warming, *J. Clim.*, *20*, 4160–4171.
- Levitus, S. (1982), Climatological atlas of the world ocean, *NOAA Prof. Pap.*, *13*, 173 pp.
- Lloyd, J., L. Park, A. Kuijpers, and M. Moros (2005), Early Holocene palaeoceanography and deglacial chronology of Disko Bugt, West Greenland, *Quat. Sci. Rev.*, *24*, 1741–1755.
- Lohmann, K., H. Drange, and M. Bentsen (2009), Response of the North Atlantic subpolar gyre to persistent North Atlantic oscillation like forcing, *Clim. Dyn.*, *32*, 273–285.
- Mignot, J., A. Levermann, and A. Griesel (2006), A decomposition of the Atlantic meridional overturning circulation into physical components using its sensitivity to vertical diffusivity, *J. Phys. Oceanogr.*, *36*, 636–650.
- Montoya, M., A. Griesel, A. Levermann, J. Mignot, M. Hofmann, A. Ganopolski, and S. Rahmstorf (2005), The Earth system model of intermediate complexity CLIMBER-3 α . Part I: Description and performance for present-day conditions, *Clim. Dyn.*, *25*, 237–263.
- Montoya, M., A. Born, and A. Levermann (2010), Reversed North Atlantic gyre dynamics in present and glacial climates, *Clim. Dyn.*, doi:10.1007/s00382-009-0729-y, in press.
- Oppo, D. W., J. F. McManus, and J. L. Cullen (2003), Deep-water variability in the Holocene epoch, *Nature*, *422*, 277–278.
- Raynaud, D., J.-M. Barnola, J. Chappellaz, T. Blunier, A. Indermühle, and B. Stauffer (2000), The ice record of greenhouse gases: A view in the context of future changes, *Quat. Sci. Rev.*, *19*, 9–17.
- Read, J. (2001), Water masses and circulation of the northeast Atlantic subpolar gyre, *Prog. Oceanogr.*, *48*, 461–510.
- Ren, J., H. Jiang, M.-S. Seidenkrantz, and A. Kuijpers (2009), A diatom-based reconstruction of Early Holocene hydrographic and climatic change in a southwest Greenland fjord, *Mar. Micropaleontol.*, *70*, 166–176.
- Renssen, H., H. Goosse, and T. Fichefet (2005), Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic seas during the Holocene, *Geophys. Res. Lett.*, *32*, L08711, doi:10.1029/2005GL022462.
- Sachs, J. P. (2007), Cooling of Northwest Atlantic slope waters during the Holocene, *Geophys. Res. Lett.*, *34*, L03609, doi:10.1029/2006GL028495.
- Solignac, S., A. de Vernal, and C. Hillaire-Marcel (2004), Holocene sea-surface conditions in the North Atlantic—Contrasted trends and regimes in the western and eastern sectors (Labrador Sea vs. Iceland Basin), *Quat. Sci. Rev.*, *23*, 319–334.
- Spall, M. A. (2005), Buoyancy-forced circulations in shallow marginal seas, *J. Mar. Res.*, *63*, 729–752.
- Thornalley, D. J. R., H. Elderfield, and I. N. McCave (2009), Holocene oscillations in temperature and salinity of the surface North Atlantic, *Nature*, *457*, 711–714.
- Treguier, A. M., S. Theetten, E. P. Chassignet, T. Penduff, R. Smith, L. Talley, J. O. Beismann, and C. Böning (2005), The North Atlantic Subpolar Gyre in four high-resolution models, *J. Phys. Oceanogr.*, *35*, 757–774.
- Wiersma, A. P., H. Renssen, H. Goosse, and T. Fichefet (2006), Evaluation of different freshwater forcing scenarios for the 8.2 ka BP event in a coupled climate model, *Clim. Dyn.*, *27*, 831–849.