

The 8.2k event: abrupt transition of the subpolar gyre towards a modern North Atlantic circulation

A. Born,^{1,2} and A. Levermann,^{3,4}

Abstract. Comparing coupled climate model simulations with marine proxy data, we find an abrupt transition of the Atlantic subpolar gyre and thereby link two major North Atlantic climate events of the early Holocene: the 8.2k event and the onset of Labrador Sea water formation.

The 8.2k event is the largest climatic signal of our present interglacial with a widespread cooling in the North Atlantic region about 8200 years before present. It coincides with a meltwater outburst from North American proglacial lakes. The associated freshwater flood is believed to have weakened the Atlantic meridional overturning circulation and northward heat transport, followed by a recovery of the deep ocean circulation and rising temperatures after a few centuries. Marine paleo records of the Labrador Sea, however, date the onset of deep water formation in Labrador Sea to the same time. The subsequent strengthening of the slope current system created a regional signal recorded as an abrupt and persistent surface temperature decrease in a number of sediment cores. Although similarities in timing are compelling, a mechanism that resolves the apparent contradiction of enhanced sinking in response to increased buoyancy from the lake drainage was missing. Our simulations suggest that the transition was due to an abrupt and persistent strengthening of the Atlantic subpolar gyre. The intense freshwater pulse triggered a transition of the gyre circulation into a different mode of operation, stabilized by internal feedbacks and persistent after the cessation of the perturbation. As a direct consequence, deep water formation around its center was intensified. This corresponds to the modern flow regime and stabilizes the meridional overturning circulation, possibly contributing to the Holocene's climatic stability.

1. Introduction

During the relatively stable conditions of our present interglacial, the 8.2k event is the largest climatic signal with a widespread cooling in the North Atlantic region about 8200 years before present. It coincides with a meltwater outburst from North American proglacial lakes (*Alley and Áugústsdóttir* [2005]; *Alley et al.* [1997] and references therein). In current understanding, this caused a weakening of the Atlantic meridional overturning circulation (AMOC) and a subsequent reduction in northward heat transport, followed by a recovery of the deep ocean circulation and rising temperatures after a few centuries [*Kleiven et al.*, 2008; *Ellison et al.*, 2006; *Wiersma et al.*, 2006; *Bauer et al.*, 2004; *Hall et al.*, 2004].

This two-dimensional explanation, however, is not able to explain a number of marine paleo records which clearly call for a three-dimensional mechanism. An abrupt but persistent surface temperature decrease was reported from the western North Atlantic at the time of the 8.2k event (Fig. 2) [*Sachs*, 2007; *de Vernal and Hillaire-Marcel*, 2006; *Solignac et al.*, 2004]. It has been suggested to be associated with the onset of deep water formation in Labrador Sea and a subsequent strengthening of the slope current system, the western

branch of the subpolar gyre (SPG) [*Sachs*, 2007; *Hillaire-Marcel et al.*, 2001]. Proxy data from Reykjanes Ridge shows a similarly abrupt and persistent warming which provides evidence for an enhanced Irminger Current (Fig. 2) [*Came et al.*, 2007; *Andersen et al.*, 2004]. This is the north-eastern branch of the SPG and thus corroborates the hypothesis of a stronger gyre and enhanced convection in its center.

The concurrence of these two events raises the question of how convection can increase at the time of the most severe freshwater flood of the past 10,000 years. We argue, based on coupled climate model experiments, that this is no contradiction but that a causal relationship connects the two events. The evaluation of the underlying physical mechanism, positive feedbacks leading to a strengthening of the SPG and convection, establishes a connection between different geological proxy data and the well assessed physical understanding of the dynamical system [*Montoya et al.*, 2010; *Born et al.*, 2010, 2009; *Levermann and Born*, 2007; *Hátún et al.*, 2005; *Treguier et al.*, 2005; *Eden and Willebrand*, 2001].

2. Model description

To investigate the role of the SPG during the 8.2k event we make use of the coupled climate model CLIMBER-3 α , which comprises atmosphere and sea ice components and the oceanic general circulation model, MOM-3 (*Montoya et al.* [2005] and Supplement). It has been used in a number of sensitivity studies with glacial [*Montoya and Levermann*, 2008] and present-day boundary conditions *Mignot et al.* [2006] as well as under global warming [*Levermann et al.*, 2007]. For the present study, the model was initialized with climatological hydrography [*Levitus*, 1982] and orbital parameters for 8,200 years before present [*Berger*, 1978]. Atmospheric CO_2 concentration was set to 260 ppm [*Raynaud*

¹Bjerknes center for Climate Research, Bergen, Norway

²Geophysical Institute, University of Bergen, Bergen, Norway

³Potsdam Institute for Climate Impact Research, Potsdam, Germany

⁴Institute of Physics, Potsdam University, Potsdam, Germany




Figure 1. Temporal evolution of key quantities of the transition towards a stronger SPG during the 8.2k event. The vertical line indicates the timing of the lake Agassiz drainage, data is filtered with a 25-year running mean. **a)** Volume and heat transport of the SPG; **b)** sea surface temperature in the north-eastern (black) and south-western (red) subpolar region; **c)** maximum winter mixed layer depth in the center of the SPG and surface density in the center of the SPG (see Fig. S7). The stronger heat transport leads to a warming of the northeastern gyre region while the southwestern part cools rapidly. Surface density and mixed layer depth increase simultaneously.

et al., 2000]. The continental watershed over North America was shifted to the west and south in order to take into account changes in surface gradient due to the isostatic depression of glacial ice sheets. After the model was run into equilibrium over 2700 years, the simulated SPG exhibits a stable but weak volume transport of 19.5 Sv ($1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$). In order to simulate the lake drainage, $160 \cdot 10^{12} \text{ m}^3$ of freshwater were added to the surface of the Labrador Sea coast during a period of two years, west of the deep convection region. This value is equivalent to a two-year volume flux of 2.6 Sv, based on a recent estimate of the lake volume [Leverington *et al.*, 2002], and has been used in previous model studies [LeGrande *et al.*, 2006; Wiersma *et al.*, 2006; Bauer *et al.*, 2004]. For diagnostic purposes, a passive tracer was released simultaneously with the freshwater flux at the same location in order to track advection of the perturbation. This tracer does not influence the circulation.

3. Model results and interpretation of proxy data

In response to the meltwater release, the SPG switches into a significantly stronger mode with 30 Sv volume transport (Fig. 1a). This represents a strengthening of about 50% and is within the uncertainty of present day observations [Read, 2001; Bacon, 1997]. Further integration shows that this stronger state is stable and not only a temporary response to the freshwater pulse.

The abrupt transition is due to two positive feedbacks inherent to the SPG [Levermann and Born, 2007]. First, a stronger SPG transports less tropical saline water into the




Figure 2. Map of the subpolar North Atlantic showing the location of core sites and of major ocean currents mentioned in the text. Blue dots denote cores that show an abrupt and persistent cooling after 8 ka before present, warming is reported from the Reykjanes Ridge and marked red. Grey dots show where the 8.2k event is evident as a temporal reduction in deep current flow speed. Black arrows illustrate the surface currents (IC, Irminger Current; EGC, East Greenland Current; LC, Labrador Current; NAC, North Atlantic Current), grey arrows the Deep Western Boundary Current. For a list of full core names and references, please refer to table S1.




Figure 3. Distribution of the meltwater pulse 30 years after its release into the Labrador Sea, integrated over depth (in m). Highest concentrations are found in the Nordic Seas. Inset: Fraction of the pulse in the Nordic Seas and the Arctic ocean. After approximately 30 years, 25% of the meltwater is advected north of the Greenland Scotland ridge, followed by a decline as it is diluted globally.

Nordic Seas but accumulates these in the subpolar North Atlantic, making the center of the SPG more saline (Fig. S7 and S9). This is consistent with previous conclusions which are based on simulations with a high-resolution oceanic general circulation model and confirmed by observations [Hátún *et al.*, 2005]. Secondly, a stronger SPG results in enhanced outcropping of isopycnals and hence a more efficient removal of heat from the gyre's center by isopycnal mixing. Both effects increase the core density of the gyre compared to the relatively light exterior, sea surface drops and the corresponding geostrophic response strengthens the cyclonic SPG circulation. In addition to these self-sustaining internal feedbacks, there exists an interaction with the flow over the Greenland Scotland ridge. The meltwater perturbation reduces sinking in the Nordic Seas and therewith the supply of dense overflow waters to the northern rim of the SPG. Consequently, the outer rim of the gyre gets lighter and the gyre slightly intensifies. This triggers the two internal feedbacks mentioned above and yields the stronger SPG state (Fig. S2). Elsewhere, we have shown that a short reduction of dense deep outflow from the Nordic Seas is sufficient to trigger the transition (≥ 25 years, [Levermann and Born, 2007]) and that the described internal feedbacks dominate the dynamics. This mechanism is robust with respect to model set-up and experimental design (see Supplement).

Considering this change in surface circulation allows to combine a number of proxy records into a consistent picture of the 8.2k event. Recent work found strong evidence for a reduction of dense water supply from the Nordic Seas due to the lake Agassiz drainage [Kleiven *et al.*, 2008; Ellison *et al.*, 2006; Hall *et al.*, 2004]. However, proxy data suggests that the meltwater signal did not reach the convection region at the center of the Labrador Sea but was exported along the shelf in the Labrador Current [Hillaire-Marcel *et al.*, 2007; Keigwin *et al.*, 2005]. From there a large portion moved north-eastward and was diluted by mixing with water of the North Atlantic Current before reaching the Nordic Seas and causing a moderate reduction in convection there. Our model reproduces this path qualitatively. 30 years after the lake drainage, highest concentrations of meltwater are found in the Nordic Seas where 25% of the meltwater pulse has been advected to (Fig. 3).

After the transition, the stronger SPG circulation intensifies oceanic heat transport (Fig. 1a). More warm tropical water reaches the northern SPG while more cold water is advected towards the west and south. This results in a sea surface temperature (SST) dipole (Fig. S7). The western SPG region cools abruptly (Fig. 1b, red) while rapid warming is seen in the north-eastern SPG region (Fig. 1b, black). An abrupt and persistent SST decrease coeval with the lake drainage has indeed been reported for many locations throughout Labrador Sea and downstream Labrador Current south of Newfoundland [Sachs, 2007; de Vernal and Hillaire-Marcel, 2006; Solignac *et al.*, 2004]. Consistent with our simulations this data indicates a drastic reorganization of the slope current system. The warming signal on the eastern side of the gyre is not easily detected in proxy records because the strengthening of the surface circulation also changed the position of frontal systems. However, on Reykjanes Ridge an increase and stabilization of surface temperatures has been associated with an intensification of the Irminger Current, the northern limb of the SPG (Fig. 2) [Came *et al.*, 2007; Andersen *et al.*, 2004].

Further support for a transition towards a stronger SPG following the 8.2k event comes from increased sea surface salinities throughout the subpolar region that started similarly abrupt as the temperature changes and were equally persistent, both in our model experiments (Fig. S7, S8 and S9) and proxy data [Hillaire-Marcel *et al.*, 2007; de Vernal and Hillaire-Marcel, 2006; Solignac *et al.*, 2004]. The cooling of western slope waters can thus not be attributed to more

intense transport of cold and fresh Arctic waters through the East Greenland Current. Their upstream source must be the relatively saline Irminger Current which suggests a SPG intensification. The intensification of westward salt transport results in a fresher inflow into the Nordic Seas (Fig. S7), consistent with paleo observations [Thornalley *et al.*, 2009]. This dynamical relationship has been confirmed by present day observations [Hátún *et al.*, 2005].

Proxy data of Labrador Sea convection furthermore supports a significant reorganization of the surface circulation. In our simulations, enhanced surface salinity together with subsurface cooling in the center of the SPG results in a density increase and a subsequent intensification of convection south of the Greenland Scotland ridge (Fig. 1c). These results match well with existing data of the geological record. Little or no Labrador Sea Water was produced in the early Holocene and formation intensified not long after the meltwater outburst [Hillaire-Marcel *et al.*, 2007, 2001]. Isotopic changes in Labrador Sea sediments show a persistent reorganization of the deep current system after 8,000 years before present [Fagel *et al.*, 2004] with deeper Labrador Sea Water [Gherardi *et al.*, 2009] and a permanent reduction of deep water formed in the Nordic Seas [Evans *et al.*, 2007]. This points towards a persistent reorganization of both the horizontal surface and the deep meridional circulation, as simulated here.

In contrast to significant changes in SPG strength (47%), the AMOC weakens by just 1.5 Sv (11%) in response to the meltwater pulse, significantly less than the 30-50% reduction reported from previous model studies [Wiersma *et al.*, 2006; LeGrande *et al.*, 2006; Bauer *et al.*, 2004] (Fig. 4). The strong response in earlier simulations might be due to the application of the meltwater pulse directly on the Labrador Sea convection region with dramatic consequences for SPG and




Figure 4. The Atlantic Meridional Overturning Circulation (AMOC), decomposed into deep water contributions from sinking in the Nordic Seas (blue) and south of the Greenland Scotland ridge (red). The vertical line indicates the timing of the lake Agassiz drainage and a 25-year running mean filter is applied. The AMOC weakens abruptly in response to the freshwater event and recovers gradually over approximately 100 years. While the initial weakening is caused by the reduction of deep water supply from the Nordic Seas, the recovery is due to a delayed intensification of Labrador Sea deep water formation. Both Nordic Seas and Labrador Sea downwelling change significantly and persistently.

AMOC. Such a freshwater perturbation leads to a reduction of convection in the Labrador Sea contradicting the paleo record that shows the intensification of convection in this region. Moreover, this scenario removes a significant fraction of the freshwater pulse from the surface, inconsistent with proxy data indicating its advection in the Labrador Current. Hence, a smaller fraction is advected into the Nordic Seas which is crucial to initiate the mechanism amplifying the SPG as discussed above (see Supplement for further discussion). The initial AMOC weakening in our simulation is followed by a rapid recovery over approximately 100 years and then a more gradual increase. The decomposition into deep water formation regions shows that the initial weakening is due to reduced sinking in the Nordic Seas, reaching its minimum after 150 years. The recovery and weak overall reduction is due to an increase in deep water formation south of the Greenland Scotland ridge with a time lag of approximately 50 years (Fig. 4, 1c).

A distinctive reduction in deep current flow speeds, probably delayed by at least several decades, is reported from the Greenland Scotland ridge overflows and off the southern tip of Greenland (Fig. 2) [Kleiven et al., 2008; Ellison et al., 2006; Hall et al., 2004]. This is upstream from where Labrador Sea Water joins the Deep Western Boundary Current, the southward flowing branch of the AMOC. Further south in the North Atlantic, the reduction of northern source deep waters is minor for the 8.2k event if recorded at all [Keigwin et al., 2005; Oppo et al., 2003; Keigwin and Boyle, 2000]. The onset of Labrador Sea Water formation following the meltwater pulse might explain this discrepancy.

4. Summary and Conclusions

Expanding previous studies that focused on the AMOC response, i.e. changes in the meridional circulation, to the lake Agassiz drainage, here we propose that many observed but hitherto unexplained abrupt changes and discrepancies require taking into account a reorganization of the horizontal circulation. The most striking result is the persistent strengthening of the SPG in response to the short freshwater pulse. The transition between the two circulation patterns is triggered by an external positive feedback and stabilized by two positive feedback mechanisms within the SPG.

While our results do not contradict an abrupt and considerable AMOC reduction in response to the lake Agassiz drainage, it might have been relatively short lived and weaker than suggested by previous simulations for two reasons: (1) The freshwater flood had the biggest impact not in the Labrador Sea but primarily affected deep water formation in the Nordic Seas after mixing in the North Atlantic Current (Fig. 3), and (2) this deep water reduction was partly compensated by enhanced sinking in the Labrador Sea (Fig. 1c, 4). The latter mechanism has already been observed in models and data on time scales of several millennia [Renssen et al., 2005; Solignac et al., 2004].

In our model the reorganization of the SPG surface circulation and subsequent changes in heat and salt advection provide the precondition for a more intense Labrador Sea convection and stabilize it. It has been suggested that this circulation mode is a unique feature of the Holocene, with Labrador Sea Water probably missing in the warmer climate of the last interglacial [Hillaire-Marcel et al., 2001] and implications for the stability of the AMOC by the end of this century. Our results might provide the base for a future investigations of these hypotheses.

Acknowledgments. We gratefully acknowledge useful discussions with Ulysses S. Ninnemann and Kerim H. Nisancioglu. A.B. was funded by the Marie Curie Actions project NICE

(MRTN-CT-2006-036127). This is publication no. XXXX from the Bjerknes Centre for Climate Research.

References

- Alley, R. B., and A. M. Águústsdóttir (2005), The 8k event: cause and consequences of a major Holocene abrupt climate change, *Quaternary Science Reviews*, *24*, 1123–1149.
- Alley, R. B., P. A. Mayewski, T. Sowers, M. Stuiver, K. C. Taylor, and P. U. Clark (1997), Holocene climatic instability: a prominent, widespread event 8200 years ago, *Geology*, *25*, 483–486.
- Andersen, C., N. Koc, and M. Moros (2004), A highly unstable Holocene climate in the subpolar North Atlantic: evidence from diatoms, *Quaternary Science Reviews*, *23*, 2155–2166.
- Bacon, S. (1997), Circulation and Fluxes in the North Atlantic between Greenland and Ireland, *Journal of Physical Oceanography*, *27*, 1420–1435.
- Bauer, E., A. Ganopolski, and M. Montoya (2004), Simulation of the cold climate event 8200 years ago by meltwater outburst from Lake Agassiz, *Paleoceanography*, *19*, PA3014.
- Berger, A. (1978), Long-Term Variations of Caloric Solar Radiation Resulting from the Earth’s Orbital Elements, *Quaternary Research*, *9*, 139–167.
- Born, A., A. Levermann, and J. Mignot (2009), Sensitivity of the Atlantic ocean circulation to a hydraulic overflow parameterisation in a coarse resolution model: Response of the subpolar gyre, *Ocean Modelling*, *27* (3–4), 130–142.
- Born, A., K. H. Nisancioglu, and P. Braconnot (2010), Sea ice induced changes in ocean circulation during the Eemian, *Climate Dynamics*, online, doi:10.1007/s00382-009-0709-2.
- Came, R. E., D. W. Oppo, and J. F. McManus (2007), Amplitude and timing of temperature and salinity variability in the subpolar North Atlantic over the past 10 k.y., *Geology*, *35*, 315–318.
- de Vernal, A., and C. Hillaire-Marcel (2006), Provincialism in trends and high frequency changes in the northwest North Atlantic during the Holocene, *Global and Planetary Change*, *54*, 263–290.
- Eden, C., and J. Willebrand (2001), Mechanism of Interannual to Decadal Variability of the North Atlantic Circulation, *Journal of Climate*, *14*, 2266–2280.
- Ellison, C. R. W., M. R. Chapman, and I. R. Hall (2006), Surface and Deep Ocean Interactions During the Cold Climate Event 8200 Years Ago, *Science*, *312*, 1929–1932.
- Evans, H. K., I. R. Hall, G. G. Bianchi, and D. W. Oppo (2007), Intermediate water links to Deep Western Boundary Current variability in the subtropical NW Atlantic during marine isotope stages 5 and 4, *Paleoceanography*, *22*, PA3209.
- Fagel, N., C. Hillaire-Marcel, M. Humblet, R. Brasseur, D. Weis, and R. Stevenson (2004), Nd and pb isotope signatures of the clay-size fraction of Labrador Sea sediments during the Holocene: Implications for the inception of the modern deep circulation pattern, *Paleoceanography*, *19*, PA3002.
- Gherardi, J.-M., L. Labeyrie, S. Nave, R. Francois, J. F. McManus, and E. Cortijo (2009), Glacial-interglacial circulation changes inferred from 231Pa/230Th sedimentary record in the North Atlantic region, *Paleoceanography*, *24*, PA2204.
- Hall, I. R., G. Bianchi, and J. R. Evans (2004), Centennial to millennial scale Holocene climate-deep water linkage in the North Atlantic, *Quaternary Science Reviews*, *23*, 1529–1536.
- Hátún, H., A. B. Sandø, H. Drange, B. Hansen, and H. Valdimarsson (2005), Influence of the atlantic subpolar gyre on the thermohaline circulation, *Science*, *309*, 1841–1844.
- Hillaire-Marcel, C., A. de Vernal, A. Bilodeau, and A. J. Weaver (2001), Absence of deep-water formation in the Labrador Sea during the last interglacial period, *Nature*, *410*, 1073–1077.
- Hillaire-Marcel, C., A. de Vernal, and D. J. W. Piper (2007), Lake Agassiz Final drainage event in the northwest North Atlantic, *Geophysical Research Letters*, *34*, L15,601.
- Keigwin, L. D., and E. A. Boyle (2000), Detecting holocene changes in thermohaline circulation., *Proceedings of the National Academy of Sciences (US)*, *97*, 1343–1346.
- Keigwin, L. D., J. P. Sachs, Y. Rosenthal, and E. A. Boyle (2005), The 8200 year B.P. event in the slope water system, western subpolar North Atlantic., *Paleoceanography*, *20*, PA2003.
- Kleiven, H. F., C. Kissel, C. Laj, U. S. Ninnemann, T. O. Richter, and E. Cortijo (2008), Reduced North Atlantic Deep Water Coeval with the Glacial Lake Agassiz Fresh Water Outburst., *Science*, *319*, 60–64.

- LeGrande, A. N., G. A. Schmidt, D. T. Shindell, C. V. Field, R. L. Miller, D. M. Koch, G. Faluvegi, and G. Hoffmann (2006), Consistent simulations of multiple proxy responses to an abrupt climate change event, *Proceedings of the National Academy of Sciences (US)*, *103*, 837–842.
- Leverington, D., H. D. Mann, and J. T. Teller (2002), Changes in bathymetry and volume of glacial Lake Agassiz between 9200 and 7700 14C yr BP, *Quaternary Research*, *57*, 244–252.
- Levermann, A., and A. Born (2007), Bistability of the Atlantic subpolar gyre in a coarse-resolution model, *Geophysical Research Letters*, *34*, L24,605.
- Levermann, A., J. Mignot, S. Nawrath, and S. Rahmstorf (2007), The role of northern sea ice cover for the weakening of the thermohaline circulation under global warming, *Journal of Climate*, *20*, 4160–4171.
- Levitus, S. (1982), *Climatological atlas of the world ocean. NOAA Professional Paper, vol 13*, 173 pp. pp., US Dept of Commerce, Washington DC.
- Mignot, J., A. Levermann, and A. Griesel (2006), A decomposition of the Atlantic meridional overturning circulation into physical components using its sensitivity to vertical diffusivity., *Journal of Physical Oceanography*, *36*, 636–650.
- Montoya, M., and A. Levermann (2008), Surface wind stress threshold for glacial Atlantic overturning, *Geophysical Research Letters*, *35*, L03,608.
- Montoya, M., A. Griesel, A. Levermann, J. Mignot, M. Hofmann, A. Ganopolski, and S. Rahmstorf (2005), The Earth System Model of Intermediate Complexity CLIMBER-3 α . Part I: description and performance for present-day conditions, *Climate Dynamics*, *25*, 237–263.
- Montoya, M., A. Born, and A. Levermann (2010), Reversed North Atlantic gyre dynamics in glacial climate, *Climate Dynamics*, *accepted*.
- Oppo, D. W., J. F. McManus, and J. L. Cullen (2003), Deepwater variability in the Holocene epoch, *Nature*, *422*, 277–278.
- Raynaud, D., J.-M. Barnola, J. Chappellaz, T. Blunier, A. Indermühle, and B. Stauffer (2000), The ice record of Greenhouse gases: a view in the context of future changes, *Quaternary Science Reviews*, *19*, 9–17.
- Read, J. (2001), Water masses and circulation of the northeast Atlantic subpolar gyre, *Progress in Oceanography*, *48*, 461–510.
- Renssen, H., H. Goosse, and T. Fichefet (2005), Contrasting trends in North Atlantic deep-water formation in the Labrador Sea and Nordic Seas during the Holocene, *Geophysical Research Letters*, *32*, L08,711.
- Sachs, J. P. (2007), Cooling of Northwest Atlantic slope waters during the Holocene, *Geophysical Research Letters*, *34*, L03,609.
- Solignac, S., A. de Vernal, and C. Hillaire-Marcel (2004), Holocene sea-surface conditions in the North Atlantic—contrasted trends and regimes in the western and eastern sectors (Labrador Sea vs. Iceland Basin), *Quaternary Science Reviews*, *23*, 319–334.
- Thornalley, D. J. R., H. Elderfield, and I. N. McCave (2009), Holocene oscillations in temperature and salinity of the surface North Atlantic, *Nature*, *457*, 711–714.
- Treguier, A. M., S. Theetten, E. P. Chassignet, T. Penduff, R. Smith, L. Talley, J. O. Beismann, and C. Böning (2005), The North Atlantic Subpolar Gyre in Four High-Resolution Models, *Journal of Physical Oceanography*, *35*, 757–774.
- Wiersma, A. P., H. Renssen, H. Goosse, and T. Fichefet (2006), Evaluation of different freshwater forcing scenarios for the 8.2 ka bp event in a coupled climate model, *Climate Dynamics*, *27*, 831–849.

A. Born, Allégaten 55, 5007 Bergen, Norway. (andreas.born@bjerkes.uib.no)